

Lattice QCD determination of states with spin $\frac{5}{2}$ or higher in the spectrum of nucleons

Subhasish Basak,¹ R. G. Edwards,² G. T. Fleming,³ J. Juge,⁴ A. Lichtl,⁴
C. Morningstar,⁴ D. G. Richards,² I. Sato,⁵ and S. J. Wallace⁶

¹*Department of Physics, N.N.D. College, Calcutta 700092, India*

²*Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, USA*

³*Yale University, New Haven, CT 06520, USA*

⁴*Department of Physics, Carnegie Mellon University, Pittsburgh, PA 15213, USA*

⁵*Lawrence Berkeley Laboratory, Berkeley, CA 94720, USA*

⁶*University of Maryland, College Park, MD 20742, USA*

Energies for excited isospin $\frac{1}{2}$ states that include the nucleon are computed using quenched, anisotropic lattices. Baryon interpolating field operators that are used include nonlocal operators that provide G_2 irreducible representations of the octahedral group. The decomposition of spin $\frac{5}{2}$ or higher states is realized for the first time in a lattice QCD calculation. We observe patterns of degenerate energies in the irreducible representations of the octahedral group that correspond to the subduction of the continuum spin $\frac{5}{2}$ or higher.

PACS numbers: 12.38.Gc 21.10.Dr

The theoretical determination of the spectrum of baryon resonances from the fundamental quark and gluon degrees of freedom is an important goal for lattice QCD. To date there have been studies of experimental ground state energies for different baryons [1–3] but only a few results for excited state energies have been reported. [4–9] No clear determination of states with spin $\frac{5}{2}$ or higher has been reported because nonlocal operators have not been used. In this work we find degenerate energies that occur in irreducible representations of the octahedral group corresponding to the subduction of the continuum spin $\frac{5}{2}$ or higher. This provides the first lattice QCD calculation that realizes the decomposition of spins greater than $\frac{3}{2}$.

Lattice correlation functions do not correspond to definite values of total angular momentum. However, they do correspond to definite irreducible representations (irreps) of the octahedral group when the source and sink operators transform accordingly. There are six double-valued irreps of the octahedral group: three for even-parity that are labeled with a g subscript (*gerade*) and three for odd-parity that are labeled with a u subscript (*ungerade*). They are: $G_{1g}, H_g, G_{2g}, G_{1u}, H_u$ and G_{2u} .

Continuum values of total angular momenta are realized in lattice simulations by patterns of degenerate energies in the continuum limit that match the patterns for the subduction of spin J to the double-valued irreps of the octahedral group. These patterns are shown in Table I. For example, a state in one of the G_2 irreps is a signal for the subduction of continuum spin $\frac{5}{2}$ or higher.

TABLE I: The number of occurrences of double-valued irrep Λ of the octahedral group for different values of continuum spin J .

Λ	$J = \frac{1}{2}$	$\frac{3}{2}$	$\frac{5}{2}$	$\frac{7}{2}$	$\frac{9}{2}$	$\frac{11}{2}$
G_1	1	0	0	1	1	1
H	0	1	1	1	2	2
G_2	0	0	1	1	0	1

For spin $\frac{5}{2}$, there must be partner states in the H and G_2 irreps that would be degenerate in the continuum limit. For spin $\frac{7}{2}$, there must be partner states in the G_1 , H and G_2 irreps and for spin $\frac{9}{2}$ there must be one partner in the G_1 irrep and two in the H irrep. The partner states should be degenerate in the continuum limit where lattice spacing $a \rightarrow 0$.

This paper reports on work to determine the pattern of low-lying states in the isospin $\frac{1}{2}$ spectrum. We carry out an analysis in quenched lattice QCD using a moderately large number of three-quark operators. Work is in progress to use a very large number of operators.[12] Smeared quark and gluon fields are used. [10, 11] Smearing reduces the couplings to short wavelength fluctuations of the theory and provides cleaner determinations of effective energies. This is important when a large array of interpolating field operators is used in order to implement the correlation-matrix method of Refs. [13, 14].

In order to determine suitable operators, including the nonlocal ones that are required for the G_2 irreps, we have developed sets of baryon operators that transform according to irreducible representations using an analytical method based on appropriate Clebsch-Gordan coefficients for the octahedral group.[15] An alternative, automated procedure that is developed in Ref. [16] provides very large sets of operators. Both methods provide equivalent results. In this work, we use for positive parity the three-quark operators defined in Tables VI, VII, and X of Ref. [15]. These comprise a complete set of quasiloal operators plus the simplest set of nonlocal operators that have one quark displaced relative to the other two quarks.

Negative-parity operators are obtained by applying the charge-conjugation transformation to the positive-parity operators. A three-quark operator that transforms according to irrep Λ and row λ of the octahedral group is related by charge conjugation to an operator that transforms according to irrep Λ_c and row λ_c and has opposite

parity, i.e.,

$$\mathcal{C}\overline{B}_k^{(\Lambda\lambda)}\mathcal{C}^\dagger = -B_k^{(\Lambda_c\lambda_c)}, \quad \mathcal{C}B_k^{(\Lambda\lambda)}\mathcal{C}^\dagger = -\overline{B}_k^{(\Lambda_c\lambda_c)*}. \quad (1)$$

If Λ is G_{1g}, G_{2g} or H_g , then Λ_c is G_{1u}, G_{2u} or H_u , respectively, and vice versa. The row labels are related by $\lambda_c = d_\Lambda + 1 - \lambda$, where d_Λ is the dimension of the irrep: $d_{G_1} = 2, d_H = 4, d_{G_2} = 2$.

A Hermitian matrix of correlation functions for irrep Λ and row λ is obtained by including a γ_4 matrix for each quark field in the source operator $\overline{B}_{k'}^{(\Lambda\lambda)}(\mathbf{0}, 0)$. This gives

$$C_{kk'}^{(\Lambda\lambda)}(t) = \mathcal{P}_{k'}^{(\Lambda)} \sum_{\mathbf{x}} \langle 0 | T \left(B_k^{(\Lambda\lambda)}(\mathbf{x}, t) \overline{B}_{k'}^{(\Lambda\lambda)}(\mathbf{0}, 0) \right) | 0 \rangle, \quad (2)$$

where the factor $\mathcal{P}_{k'}^{(\Lambda)}$ is ± 1 and it arises from evaluating the action of the γ_4 matrices in the source operator. Subscripts k and k' refer to different embeddings of irrep Λ . The relation between matrices of correlation functions with different parities is [16],

$$C_{kk'}^{(\Lambda\lambda)}(t) = -\eta_t C_{kk'}^{(\Lambda_c\lambda_c)*}(T - t). \quad (3)$$

where $\eta_t = 1$ for periodic boundary conditions and $\eta_t = -1$ for anti-periodic boundary conditions. Applying charge conjugation and time-reversal to a correlation function produces one for the opposite parity because of PCT symmetry. We use a sufficiently long time extent, T , so that this symmetry provides two independent correlation functions for each parity.

Analyses of excited state energies are based upon the matrices of correlation functions of Eq. (2). In order to extract the spectrum of energies from the matrix of correlation functions, we first average over rows because they provide equivalent results owing to octahedral symmetry. We then solve the following generalized eigenvalue equation,

$$\sum_{k'} \tilde{C}_{kk'}^{(\Lambda)}(t) v_{k'}^{(n)}(t, t_0) = \alpha^{(n)}(t, t_0) \sum_{k'} \tilde{C}_{kk'}^{(\Lambda)}(t_0) v_{k'}^{(n)}(t, t_0), \quad (4)$$

where superscript n labels the eigenstates. The symbol $\tilde{C}^{(\Lambda)}$ indicates that the appropriate average over rows has been performed. The reference time t_0 in Eq. (4) is taken near the source time $t = 0$ in order to have significant contributions from excited states. The principal eigenvalues $\alpha^{(n)}(t, t_0)$ are related to the energy E_n by [17]

$$\alpha^{(n)}(t, t_0) \simeq e^{-E_n(t-t_0)} \left(1 + \mathcal{O}(e^{-|\delta E|t}) \right), \quad (5)$$

where δE is the difference between E_n and the next closest energy. We have determined effective energies E_n by fitting the generalized eigenvalues to the leading term of Eq. (5).

Methods that improve the signal-to-noise ratio are very important for extracting excited state energies from lattice QCD simulations. The correlation-matrix method optimizes the operators so as to achieve early plateaus

of the effective energies and the use of PCT symmetry increases the statistics. In addition, we use anisotropic lattices with temporal lattice spacing a_t one-third of the spatial lattice spacing a_s . The finer spacing increases the number of time-slices for extraction of energies before the signal for a high-energy state decays to the level of the noise.

In this work anisotropic lattices with two different volumes are used: 239 gauge field configurations are used for a $16^3 \times 64$ lattice and 167 configurations are used for a $24^3 \times 64$ lattice. Because of the use of PCT symmetry, the size of the statistical ensemble is effectively doubled. Gauge-field configurations are generated using the anisotropic, unimproved Wilson gauge action [18–21] in the quenched approximation with $\beta = 6.1$ and the pion mass is 490 MeV. For both lattices, the temporal lattice spacing corresponds to $a_t^{-1} = 6.05(1)$ GeV [8] as determined from the string tension. Calculations are performed using the Chroma software. [22] Further details of the action and lattice parameters will be presented elsewhere.

We have extracted energies for isospin $\frac{1}{2}$ and $\frac{3}{2}$ channels by diagonalizing matrices of correlation functions formed from three-quark operators that share the same octahedral symmetry. Starting with a large number of operators, we first eliminate operators that have less influence on the determination of the effective energies and the eigenvectors $\mathbf{v}^{(n)}$ of the low-lying states. Once a set of good operators is obtained, we form final matrices of correlation functions, diagonalize them and extract effective energies. This procedure yields solid results for energies of low-lying states and is more efficient than diagonalizing matrices of the largest dimension. We have obtained 17 energies for nucleonic states and 11 energies for delta baryon states. In this paper, we focus on the nucleon channel and the determination of spin.

Our lattice results for the spectrum are shown in Fig. 1. For positive parity, the nucleon state is very well determined at lattice energy $a_t E = 0.193(3)$. It is an isolated G_{1g} state corresponding to spin $\frac{1}{2}$. Well above the nucleon is a group of three degenerate states (within errors) in the G_{1g} , H_g and G_{2g} irreps. The results on both lattice volumes are essentially the same, although the lowest G_{2g} state in the smaller volume has a larger error.

Energies of G_2 states are the most difficult to extract because the plateaus are relatively short-lived, signal-to-noise ratios are large, and the number of operators that we have used for the matrices of correlation functions is limited. The limitation on the number of operators is because we restrict the operators used in this work to ones that can be constructed from one-link displacements. In work that is in progress, more varied types of operators are used to obtain very larger numbers of G_2 operators, which is essential for determining states with higher energies.

Figure 2 shows effective masses obtained for the lowest energy G_{1g} , H_g and G_{2g} states. The horizontal lines indicate the time ranges used to fit the principal eigenvalues




FIG. 1: The energies obtained for each symmetry channel of isospin $\frac{1}{2}$ baryons are shown based on the $24^3 \times 64$ lattice data. The scale on the left side shows energies in lattice units and the scale on the right side shows energies in GeV. Errors are indicated by the vertical size of the box.




FIG. 2: Effective energies for the lowest states in G_{1g} , H_g and G_{2g} irreps. Time ranges used for and energies obtained from fits to the principal eigenvalues are shown by horizontal lines.

$\alpha^{(n)}(t, t_0)$ by a single exponential form. The resulting mean energy value and the error range are indicated by the lines.

Although there are substantial discretization errors with the quark action that is used, and they could contribute differently in the different irreps, clear patterns in the degeneracies emerge. Focusing on the group of three positive-parity states near lattice energy $a_t E = 0.36$ in Fig. 1, two interpretations are possible. A.) The group consists of a spin $\frac{1}{2}$ state and a spin $\frac{5}{2}$ state that are accidentally degenerate. In this case the G_{1g} state corresponds to spin $\frac{1}{2}$ and the H_g and G_{2g} partner states

correspond to the subduction of spin $\frac{5}{2}$. B.) The group consists of a single state with the degenerate G_{1g} , H_g and G_{2g} partner states corresponding to the subduction of spin $\frac{7}{2}$.

In the physical spectrum of positive-parity nucleon resonances, the lowest excited state, $N(1440, \frac{1}{2}^+)$, lies below all negative parity states. We do not find a signal for a positive-parity excitation that has lower energy than the negative-parity excitations at this quark mass. The next two excited nucleon states are essentially degenerate, namely, $N(1680, \frac{5}{2}^+)$ and $N(1710, \frac{1}{2}^+)$, each with a width of about 100 MeV. Spin $\frac{7}{2}$ states occur only at significantly higher energy (1990 MeV). Primarily because of the absence of spin $\frac{7}{2}$ in the low-lying spectrum, interpretation A.) of our lattice results is more consistent with the physical pattern of energies and spins. In the absence of G_2 operators, previous lattice studies have assumed that states obtained with H irrep operators correspond to continuum spin $\frac{3}{2}$. [4, 6, 7, 9]. With the G_2 operators, we find low-lying states in irrep H that are consistent with the subduction of spin $\frac{5}{2}$ or higher.

In the negative-parity spectrum, we also obtain essentially the same results for both lattice volumes. The three lowest states shown on the right half of Fig. 1 are unambiguously identified as follows: the lowest G_{1u} state corresponds to spin $\frac{1}{2}$ and the lowest two H_u states correspond to distinct spin $\frac{3}{2}$ states. Above these is a group of three states with roughly the same lattice energy: $a_t E \approx 0.33$ (within errors). Again there are two possible interpretations. C.) The group consists of a spin $\frac{1}{2}$ state in G_{1u} that is accidentally degenerate with a spin $\frac{5}{2}$ state, the latter having degenerate partner states in H_u and G_{2u} . D.) The group consists a spin $\frac{7}{2}$ state having degenerate partner states in G_{1u} , H_u and G_{2u} .

The pattern of low-lying physical states starts with $N(\frac{3}{2}^-, 1520)$, $N(\frac{1}{2}^-, 1535)$ and $N(\frac{3}{2}^-, 1700)$. These should show up as distinct H_u , G_{1u} and H_u states on the lattice, in agreement with the three lowest negative-parity states in Fig. 1. The next physical states include $N(\frac{1}{2}^-, 1680)$ and $N(\frac{5}{2}^-, 1675)$, which essentially are degenerate. They should show up as degenerate G_{1u} , H_u and G_{2u} states on the lattice. This pattern of spins is consistent with interpretation C.) of the lattice states at lattice energy $a_t E \approx 0.33$. The pattern of energies of the physical states has $N(\frac{5}{2}^-, 1675)$ a little lower in energy than $N(\frac{3}{2}^-, 1700)$, but the lattice results at lattice spacing $0.1F$ place the spin $\frac{5}{2}$ state above the spin $\frac{3}{2}$ state. Study of the continuum limit of the lattice spectrum is required in order to resolve these issues.

Because the minimum spin that is contained in the G_2 irrep is $\frac{5}{2}$, we have found strong evidence for spin $\frac{5}{2}$ or higher for both parities in our lattice spectra for isospin $\frac{1}{2}$. We also have found evidence for degenerate partner states corresponding to the subduction of spin $\frac{5}{2}$ or higher to the octahedral irreps. The lattice results for the low-lying excited states of isospin $\frac{1}{2}$ provide the

correct number of octahedral states for the subduction of the spins of the low-lying physical states. These results are significant but they are based on the quenched approximation and a 490 MeV pion mass. Similar calculations in full QCD with several pion masses and several lattice spacings are planned and work is under way to calculate the anisotropic gauge configurations.

Acknowledgments

This work was supported by the U.S. National Science Foundation under Award PHY-0354982 and by the U.S. Department of Energy under contracts DE-AC05-06OR23177 and DE-FG02-93ER-40762.

-
- [1] S. Aoki *et al.* [CP-PACS Collaboration], Phys. Rev. Lett. **84**, 238 (2000) [arXiv:hep-lat/9904012].
 - [2] C. T. H. Davies *et al.* [HPQCD Collaboration], Phys. Rev. Lett. **92**, 022001 (2004) [arXiv:hep-lat/0304004].
 - [3] A. Ali Khan *et al.* [CP-PACS Collaboration], Phys. Rev. D **65**, 054505 (2002) [Erratum-ibid. D **67**, 059901 (2003)] [arXiv:hep-lat/0105015].
 - [4] L. Zhou and F. X. Lee, arXiv:hep-lat/0604023.
 - [5] T. Burch, C. Gattringer, L. Y. Glozman, C. Hagen, D. Hierl, C. B. Lang and A. Schafer, arXiv:hep-lat/0604019.
 - [6] K. Sasaki and S. Sasaki, PoS **LAT2005**, 060 (2005) [arXiv:hep-lat/0508026].
 - [7] K. Sasaki and S. Sasaki, Phys. Rev. D **72**, 034502 (2005) [arXiv:hep-lat/0503026].
 - [8] S. Basak *et al.* [LHP Collaboration], Nucl. Phys. Proc. Suppl. **140**, 278 (2005) [arXiv:hep-lat/0409082].
 - [9] J. M. Zanotti, D. B. Leinweber, A. G. Williams, J. B. Zhang, W. Melnitchouk and S. Choe [CSSM Lattice collaboration], Phys. Rev. D **68**, 054506 (2003) [arXiv:hep-lat/0304001].
 - [10] M. Albanese *et al.* (APE Collaboration), Phys. Lett. B **192**, 163 (1987).
 - [11] M. G. Alford, T. Klassen and P. Lepage, Nucl. Phys. Proc. Suppl. **47**, 370 (1996) [arXiv:hep-lat/9509087].
 - [12] A. Lichtl, [arXiv:hep-lat/0609019].
 - [13] C. Michael, Nucl. Phys. B **259**, 58 (1985).
 - [14] M. Lüscher and U. Wolff, Nucl. Phys. B **339**, 222 (1990).
 - [15] S. Basak *et al.* [Lattice Hadron Physics Collaboration (LHPC)], Phys. Rev. D **72**, 074501 (2005) [arXiv:hep-lat/0508018].
 - [16] S. Basak *et al.*, Phys. Rev. D **72**, 094506 (2005) [arXiv:hep-lat/0506029].
 - [17] M. Luscher and U. Wolff, Nucl. Phys. B **339**, 222 (1990).
 - [18] P. Chen, Phys. Rev. D **64**, 034509 (2001) [arXiv:hep-lat/0006019].
 - [19] T. R. Klassen, Nucl. Phys. B **533**, 557 (1998) [arXiv:hep-lat/9803010].
 - [20] F. Karsch, J. Engels and T. Scheideler, Nucl. Phys. Proc. Suppl. **63**, 427 (1998) [arXiv:hep-lat/9709011].
 - [21] M. Fujisaki *et al.* [QCD-TARO Collaboration], Nucl. Phys. Proc. Suppl. **53**, 426 (1997) [arXiv:hep-lat/9609021].
 - [22] R. G. Edwards (LHPC Collaboration) and B. Joo (UKQCD Collaboration), "The Chroma Software System for Lattice QCD", arXiv:hep-lat/0409003, Proceedings of the 22nd International Symposium for Lattice Field Theory (Lattice2004), Nucl. Phys B140 (Proc. Suppl) p832, 2005.